arXiv:alg-geom/9707018v2 11 Aug 1997

ON SIMPLICIAL COMMUTATIVE RINGS WITH VANISHING ANDRÉ-QUILLEN HOMOLOGY

JAMES M. TURNER

Abstract. We propose a generalization of a conjecture of D. Quillen, on the vanishing of André-Quillen homology, to simplicial commutative rings. This conjecture characterizes a notion of local complete intersection, extended to the simplicial setting, under a suitable hypothesis on the local characteristic. Further, under the condition of finite-type homology, we then prove the conjecture in the case of a simplicial commutative algebra augmented over a field of non-zero characteristic. As a consequence, we obtain a proof of Quillen's conjecture for a Noetherian commutative algebra - again augmented over a field of non-zero characteristic.

1. Introduction

In [1] and [12], M. André and D. Quillen constructed the notion of a homology $D_*(A|R;M)$ for a commutative algebra A, over a ring R, and an A-module M. It was then conjectured (see Section 5 of [12]) that, under suitable conditions on R and A, the vanishing of the homology in sufficiently high degrees determines A as a local complete intersection. In particular, for local rings, the conjecture takes the following form.

Conjecture 1.1. *Let R be a (Noetherian, commutative) local ring with residue field $\mathbb{F}$, and let $D_s(\mathbb{F}|R) = D_s(\mathbb{F}|R;\mathbb{F})$, $s \geq 0$. Then the following are equivalent:*

1. $D_s(\mathbb{F}|R) = 0, \quad s >> 0$;
2. $D_s(\mathbb{F}|R) = 0, \quad s \geq 3$;
3. *R is a complete intersection.*

In this form, 1.1 was proven by L. Avramov, in outline form, in [2], and, in much greater generality, in [3]. Recall that a local ring R is a complete intersection if its I-adic completion $\hat{R}$ is a quotient of a complete regular ring by an ideal generated by a regular sequence. From this description and the properties of André-Quillen homology, the implications (3) ⇒ (2) ⇒ (1) in 1.1 are immediate.

The objective of this paper is to extend a version of 1.1 for simplicial local rings in an effort to bring the full power of simplicial homotopy theory to bear on this type of problem and thereby obtain a different proof of 1.1 closer in spirit to the topological results of J.-P. Serre in [14] and Y. Umeda in [16].

Date: August, 1997.

1991 *Mathematics Subject Classification.* Primary: 13D03, 18G30, 18G55; Secondary: 13D40, 18G20, 55S45, 55T10.

Key words and phrases. simplicial commutative algebras, André-Quillen homology, complete intersections, Serre spectral sequences, simplicial dimension, Postnikov envelopes, Poincaré series.

Research was partially supported by an NSF-NATO postdoctoral fellowship.

In [13], D. Quillen gave a construction of André-Quillen homology $D_*(A|B;M)$ where B is a simplicial commutative ring, A a simplicial commutative B-algebra, and M a simplicial A-module.

Let $\mathcal{R}_{\mathbb{F}}$ be the category of (commutative) local rings, with residue field $\mathbb{F}$, and $s\mathcal{R}_{\mathbb{F}}$ the category of simplicial objects over $\mathcal{R}_{\mathbb{F}}$. It follows from [11] that $s\mathcal{R}_{\mathbb{F}}$ has three classes of maps, called weak equivalences, fibrations, and cofibrations, giving it the structure of a closed simplicial model category. Using this structure, we say that a simplicial local ring R is an *n-extension* if there is a cofibration sequence

$$\Sigma^{n-1}S_0 \to R \to S_1,$$

in the homotopy category $Ho(s\mathcal{R}_{\mathbb{F}})$, such that S_0 is polynomial in $\mathcal{R}_{\mathbb{F}}$ and $\hat{S}_1 \cong \Sigma^n \bar{S}_1$ in $Ho(s\mathcal{R}_{\mathbb{F}})$. Here Σ denotes the suspension in $Ho(s\mathcal{R}_{\mathbb{F}})$. (See §I.2 and §I.3 of [11] for the theory of suspension and cofibration sequences in homotopical algebra.)

Definition 1.2. *Let R be an object of $s\mathcal{R}_{\mathbb{F}}$. Then:*

1. *R is* **regular** *if R is a 1-extension with $\bar{S}_1$ smooth in $\mathcal{R}_{\mathbb{F}}$.*
2. *R is a* **complete intersection** *if it is a 1-extension with $\bar{S}_1$ regular in $s\mathcal{R}_{\mathbb{F}}$.*
3. *R is Q-***bounded** *if $(Q\pi_*R)_s = 0$ for $s >> 0$, and* bounded *if $\pi_s R = 0$ for $s \gg 0$.*
4. *The* **simplicial dimension** *of R is the integer*
$$s\cdot\dim R = \max\{s|\, D_s(\mathbb{F}|R) \neq 0\}.$$
We then say that R has **finite simplicial dimension** *if $s\cdot\dim R < \infty$.*
5. *R is said to have* **finite-type homology** *provided each $D_q(\mathbb{F}|R)$ is a finite dimensional $\mathbb{F}$-vector space.*
6. *If R has both finite-type homology and finite simplicial dimension, we call R* **finite**.

Given a simplicial commutative ring R, then R is said to be locally *of any one of (1) – (6) provided $R_\wp$ is such, for each simplicial prime ideal $\wp$ in R. Given a simplicial prime ideal $\wp$ in R, we denote by $\mathbb{F}(\wp)$ the residue field of $R_\wp$ and we say that R is* locally of non-zero characteristic *provided char$\mathbb{F}(\wp) \neq 0$ for all such $\wp$.*

We can now state our proposed simplicial generalization of Conjecture 1.1.

Vanishing Conjecture 1.3. *Let R be a locally finite simplicial commutative ring which is locally of non-zero characteristic. Then R is a locally complete intersection if and only if R is locally Q-bounded.*

In the rational case, while a complete intersection may be both Q-bounded and of finite simplicial dimension, the converse is not true. See the note following Proposition 3.4 regarding counter-examples.

To demonstrate the validity of 1.3, we consider the subcategory, $\mathcal{A}_{\mathbb{F}}$, of $\mathcal{R}_{\mathbb{F}}$, consisting of augmented $\mathbb{F}$-algebras, i.e., unitary $\mathbb{F}$-algebras A together with a fixed $\mathbb{F}$-algebra map $A \to \mathbb{F}$, called the augmentation of A.

In this paper, we give evidence for Vanishing Conjecture 1.3 by proving:

Theorem 1.4. *Suppose A is a finite simplicial augmented commutative $\mathbb{F}$-algebra with char$\mathbb{F} > 0$. Then A is bounded if and only if A is a complete intersection.*

Corollary 1.5. *Let A be an augmented Noetherian commutative $\mathbb{F}$-algebra, char$\mathbb{F} > 0$. Then A is a complete intersection, as a local algebra, if and only if A has finite simplicial dimension.*

Proof. By IV.55 of [1], $H_*^Q(A)$ is of finite-type. Thus, by Theorem 1.4, A has finite simplicial dimension if and only if A is a complete intersection, as a simplicial algebra, if and only if A is a complete intersection, as a local algebra, by the classical implication of (2) ⇒ (3) in Conjecture 1.1 (see Proposition 26 of [1]). □

Organization of this paper. In this section, we review the closed simplicial model category structure for $s\mathcal{A}_{\mathbb{F}}$ and the construction and properties of homotopy and André-Quillen homology. In Section 3, we describe the notion Postnikov envelopes for objects of $s\mathcal{A}_{\mathbb{F}}$ and explore its properties. In Section 4, we study the homotopy of n-extensions. Finally, in Section 5, we introduce and study the notion of a Poincaré series for a simplicial algebra, obtaining just enough information to prove Theorem 1.4.

Acknowledgements. The author would like to thank Haynes Miller, for suggesting this project along with the direction it should take, and Jean Lannes for many useful directions as well as for making his stay in France worthwhile. Most of the work on this project was done while the author was visiting the Institut des Hautes Études Scientifique and the Ecole Polytechnique. He would like to thank them for their hospitality and use of their facilities during his stay. Finally, the author would like to thank Julie Riddleburger for putting this paper into LaTeX form.

2. Homotopy Theory of Simplicial Augmented Commutative Algebras

We now review the closed simplicial model category structure for $s\mathcal{A}_{\mathbb{F}}$. We will assume the reader is familiar with the general theory of homotopical algebra given in [11].

We call a map $f: A \to B$ in $s\mathcal{A}_{\mathbb{F}}$ a

(i) weak equivalence ($\xrightarrow{\sim}$) $\iff \pi_* f$ is an isomorphism;
(ii) fibration ($\twoheadrightarrow$) $\iff f$ surjects in positive degrees;
(iii) cofibration($\hookrightarrow$) $\iff f$ is a retract of an almost free map.

Here a map $f: A \to B$ in $s\mathcal{A}_{\mathbb{F}}$ is *almost free* if there is an almost simplicial $\mathbb{F}$-vector space (no d_0) V (see [7]) together with a map of almost simplicial $\mathbb{F}$-vector spaces $V \to IB$ such that the induced map $A \otimes S(V) \xrightarrow{\cong} B$ is an isomorphism of almost simplicial algebras. Here S is the symmetric algebra functor.

Now given a finite simplicial set K and a simplicial algebra A, define $A \wedge K$ and A^K by

$$(A \wedge K)_n = \bigotimes_{K_n} A_n$$

and

$$(A^K)_n = \prod_{K_n} A_n.$$

Here the tensor product $\otimes$ is the coproduct in $s\mathcal{A}_{\mathbb{F}}$. The product in $s\mathcal{A}_{\mathbb{F}}$ is defined as $\Lambda \times_{\mathbb{F}} \Gamma$, for Λ, Γ in $s\mathcal{A}_{\mathbb{F}}$, so that the diagram

$$\begin{tikzcd}
\Lambda \times_{\mathbb{F}} \Gamma \arrow[r] \arrow[d] & \Gamma \arrow[d, "\epsilon"] \\
\Lambda \arrow[r, "\epsilon"'] & \mathbb{F}
\end{tikzcd}$$

is a pullback of simplicial vector spaces.

Theorem 2.1. *([11], [10], and [7]) With these definitions, $s\mathcal{A}_{\mathbb{F}}$ is a closed simplicial model category.*

Given a simplicial vector space V, define its normalized chain complex NV by

$$N_nV = V_n/(\mathrm{Im}s_0 + \cdots + \mathrm{Im}s_n) \tag{2.1}$$

and $\partial : \; N_nV \to N_{n-1}V$ is $\partial = \sum_{i=0}^{n}(-1)^i d_i$. The homotopy groups π_*V of V is defined as

$$\pi_nV = H_n(NV), \quad n \geq 0.$$

Thus for A in $s\mathcal{A}_{\mathbb{F}}$ we define π_*A as above. The Eilenberg-Zilber theorem (see [9]) shows that the algebra structure on A induces an algebra structure on π_*A.

If we let $\mathcal{V}$ be the category of $\mathbb{F}$-vector spaces, then there is an adjoint pair

$$S : \; \mathcal{V} \Longleftrightarrow \mathcal{A}_{\mathbb{F}} : \; I,$$

where I is the augmentation ideal function and S is the symmetric algebra functor. For an object V in $\mathcal{V}$ and $n \geq 0$, let $K(V,n)$ be the associated Eilenberg-MacLane object in $s\mathcal{V}$ so that

$$\pi_sK(V,n) = \begin{cases} V & s = n; \\ 0 & s \neq n. \end{cases}$$

Let $S(V,n) = S(K(V,n))$, which is an object of $s\mathcal{A}_{\mathbb{F}}$.

For A in $\mathcal{A}_{\mathbb{F}}$, the indecomposable functor $QA = I(A)/I^2(A)$ which is an object of $\mathcal{V}$. Furthermore, we have an adjoint pair

$$Q : \; \mathcal{A}_{\mathbb{F}} \Longleftrightarrow \mathcal{V} : \; (-)_+$$

where V_+, for V in $\mathcal{V}$, is the simplicial algebra $V \oplus \mathbb{F}$ where

$$(v,r) \cdot (w,s) = (sv + rw, rs)$$

for $(v,r),(w,s) \in V \oplus \mathbb{F}$. $(-)_+, Q$ provides an equivalence between $\mathcal{V}$ and the category of abelian group objects in $\mathcal{A}_{\mathbb{F}}$.

For A in $s\mathcal{A}_{\mathbb{F}}$, we define its André-Quillen homology, as per [7] and [8], by

$$H_s^Q(A) = \pi_sQX, \quad s \geq 0,$$

where we choose a factorization

$$\mathbb{F} \hookrightarrow X \xrightarrow{\sim} A$$

of the unit $\mathbb{F} \to A$ as a cofibration and a trivial fibration. This definition is independent of the choice of factorization as any two are homotopic over A (note that every object of $s\mathcal{A}_{\mathbb{F}}$ is fibrant). It is known (see, for example, [10]) that

$$H_s^Q(A) = D_s(A|\mathbb{F};\mathbb{F}).$$

From the transitivity sequence, one can easily check that $D_0(\mathbb{F}|A) = 0$, and $D_{s+1}(\mathbb{F}|A) \cong H_s^Q(A)$ for all $s \geq 0$.

Now, as shown in [8],

$$\pi_n A = [S(n), A],$$

where $S(n) = S(\mathbb{F}, n)$ and $[\quad,\quad]$ denotes the morphisms in $Ho(s\mathcal{A}_{\mathbb{F}})$. Thus the primary operational structure for the homotopy groups in $s\mathcal{A}_{\mathbb{F}}$ is determined by $\pi_* S(V_0)$ for any V_0 in $s\mathcal{V}$. By Dold's theorem [6] there is a triple $\mathcal{S}$ on graded vector spaces so that

$$\pi_* S(V) \cong \mathcal{S}(\pi_* V) \tag{2.2}$$

encoding this structure. If char$\mathbb{F} = 0$, $\mathcal{S}$ is the free skew symmetric functor and, if char$\mathbb{F} > 0$, $\mathcal{S}$ is the free divided power algebra on the underlying vector space of a certain free algebra constructed from the input (see, for example, [4] and [7]).

Now recall that maps $A \xrightarrow{f} B \xrightarrow{g} C$ is a *cofibration sequence* in $Ho(s\mathcal{A}_{\mathbb{F}})$ if f is isomorphic to a cofibration $X \xrightarrow{u} Y$, of cofibrant objects, with cofibre $Y \xrightarrow{v} Z$ isomorphic to g. Thus given any map $f:\ A \to B$ in $s\mathcal{A}_{\mathbb{F}}$ there is a cofibration sequence $A \xrightarrow{f} B \to M(f)$ in $Ho(s\mathcal{A}_{\mathbb{F}})$ formed by factoring $\mathbb{F} \to A$ into $\mathbb{F} \hookrightarrow \bar{A} \overset{\sim}{\twoheadrightarrow} A$, form the diagram

$$\begin{tikzcd}
\bar{A} \arrow[r, hook] \arrow[d, two heads, "s"'] & X \arrow[d, two heads, "\wr"] \\
A \arrow[r, "f"] & B
\end{tikzcd}$$

and then let $M(f) = X \otimes_{\bar{A}} \mathbb{F}$, which is cofibrant. As an example, the *suspension* ΣA of an object A in $s\mathcal{A}_{\mathbb{F}}$ by $M(\epsilon)$, where $\epsilon:\ A \to \mathbb{F}$ is the augmentation.

Finally, recall that the *completion* $\hat{A}$ of a simplicial augmented algebra A is defined as

$$\hat{A} = lim_t A/I^t.$$

If $f: A \to B$ is a map of simplicial algebras, we denote by $\hat{f}: \hat{A} \to \hat{B}$ the induced map of completions.

We can now summarize methods for computing homotopy and André-Quillen homology that we will need for this paper.

Proposition 2.2. (1) *If $f:\ A \xrightarrow{\sim} B$ is a weak equivalence in $s\mathcal{A}_{\mathbb{F}}$, then $H_*^Q(f):\ H_*^Q(A) \xrightarrow{\cong} H_*^Q(B)$ is an isomorphism.*

(2) *for any A in $s\mathcal{A}_{\mathbb{F}}$ there is a spectral sequence*

$$E^1_{s,t} = \mathcal{S}_s(H_*^Q(A)) \Rightarrow \pi_t \hat{A},$$

called Quillen's Fundamental spectral sequence (see [12] *and* [13]*), which converges when $H_0^Q(A) = 0$.*

(3) *There is a Hurewicz homomorphism* $h:\ I\pi_*A \to H_*^Q(A)$ *such that if* A *is connected and* $H_s^Q(A)=0$ *for* $s<n$ *then* A *is* n*-connected and* $h:\ \pi_nA \xrightarrow{\cong} H_n^Q(A)$ *is an isomorphism.*

Proof. (1) is a standard result. See, for example, [12] or [8]. For (2), see chapter IV of [7] and [15]. Finally, (3) is in [7]. □

The following is a selection of results from [15].

Proposition 2.3. *Let* A *and* B *be in* $s\mathcal{A}_{\mathbb{F}}$. *Then*

1. *if* $f:A\to B$ *is an* H_*^Q*-isomorphism then* $\hat{f}:\hat{A}\to\hat{B}$ *is a weak equivalence,*
2. *if* A *is connected then* $\pi_*\hat{A}\cong\pi_*A$,
3. $H_*^Q(\hat{A})\cong H_*^Q(A)$, *and*
4. $Q\pi_*\hat{A}\cong Q\pi_*A$.

Remark. If $H_0^Q(A)=0$ then Proposition 2.3 (4) follows from a Quillen fundamental spectral sequence argument. This is due to the fact that while this spectral sequence doesn't directly converge to π_*A it does allow, under the above condition on H_0^Q, sufficient information to be extracted about the indecomposables (see [15] for further details). This case is sufficient for our needs.

Proposition 2.4. *Let* $A \xrightarrow{f} B \xrightarrow{g} C$ *be a cofibration sequence in* $Ho(s\mathcal{A}_{\mathbb{F}})$. *Then:*

1. *There is a long exact sequence*

$$\cdots \to H_{s+1}^Q(C) \xrightarrow{\partial} H_s^Q(A) \xrightarrow{H_*^Q(f)} H_s^Q(B)$$

$$\xrightarrow{H_*^Q(g)} H_s^Q(C) \xrightarrow{\partial} H_{s-1}^Q(C) \to \cdots$$

2. *There is a first quadrant spectral sequence of algebras*

$$E_{s,t}^2 = Tor_s^{\pi_*A}(\pi_*B,\mathbb{F})_t \Rightarrow \pi_{s+t}C,$$

which we refer to as the Eilenberg-Moore spectral sequence.

3. *If* A *is connected, there is a first quadrant spectral sequence of algebras*

$$E_{s,t}^2 = \pi_s(C\otimes\pi_tA) \Rightarrow \pi_{s+t}B,$$

which we refer to as the Serre spectral sequence.

4. *If* A *is connected and* C *is* n*-connected, then there is a homomorphism* $\tau:\ \pi_{n+1}C\to\pi_nA$, *called the* transgression, *such that the diagram*

$$\begin{tikzcd}
\pi_{n+1}B \arrow[r, "\pi_*f"] \arrow[d, "h"'] & \pi_{n+1}C \arrow[r, "\tau"] \arrow[d, "h"'] & \pi_nA \arrow[r, "\pi_*f"] \arrow[d, "h"] & \pi_nB \arrow[d, "h"] \\
H_{n+1}^QB \arrow[r, "H_*^Qf"] & H_{n+1}^Q(C) \arrow[r, "\partial"] & H_n^Q(A) \arrow[r, "H_*^Qf"] & H_n^QB
\end{tikzcd}$$

commutes and the top sequence is exact.

Proof. (1) is just the transitivity sequence for H^Q_*. See [7].

(2) is the spectral sequence of Theorem 6(b) in §II.6 of [11]. See also [7]. By Theorem 6(d) in §II.6 of [11], there is a 1^{st}-quadrant spectral sequence

$$E^2_{*,*} = \pi_*(B \otimes_A \pi_* A) \Rightarrow \pi_* B,$$

where $\pi_* A$ is an A-module via the augmentation $A \to \pi_0 A$. Here we can assume our cofibration sequence is a cofibration with cofibre C. Since A is connected, then $B \otimes_A \pi_* A \cong C \otimes \pi_* A$. The algebra structure follows from the construction of the spectral sequence and the fact that $A \xrightarrow{f} B$ is a map of simplicial algebras. This gives us (3).

For (4), since A is connected and C is n-connected, then in the Serre spectral sequence

$$d^{n+1}: \ \pi_{n+1} C \cong E^{n+1}_{n+1,0} \to E^{n+1}_{0,n} \cong \pi_n A,$$

which we propose is our desired map τ. From this same spectral sequence, we have $\pi_s A \cong \pi_s B$, $s < n$, and, using methods modified from the next section, we can assume that $N_s IC = 0$ for $s \leq n$ and $N_{n+1} B \twoheadrightarrow N_{n+1} C$ is surjective. Since $E^1_{n+1,0} = N_{n+1} C$ and $E^2_{n+1,0} \cong E^{n+1}_{n+1,0} \cong \pi_{n+1} C$, then d^{n+1} is constructable in precisely the same way as the boundary map in homological algebra. Since we can assume cofibrancy of our objects under consideration, then the diagram

$$\begin{tikzcd}
H_{n+1}(NB) \arrow[r] \arrow[d] & H_{n+1}(NC) \arrow[r, "d^{n+1}"] \arrow[d] & H_n(NA) \arrow[r] \arrow[d] & H_n(NB) \arrow[d] \\
H_{n+1}(NQB) \arrow[r] & H_{n+1}(NQC) \arrow[r, "\partial"] & H_n(NQA) \arrow[r] & H_n(NQB)
\end{tikzcd}$$

commutes by naturality. The result follows. □

3. Postnikov Envelopes

In this section, we construct and determine some properties of a useful tool for studying simplicial algebras. First, we recall the following standard result which will be useful for us (see section II.4 of [11]).

Lemma 3.1. *Let V and W be simplicial vector spaces. Then the map*

$$[V, W] \to Hom_{\mathcal{V}_*}(\pi_* V, \pi_* W)$$

is an isomorphism.

Proposition 3.2. *Let A in $s\mathcal{A}_{\mathbb{F}}$. Then*

1. *There is a map of simplicial algebras*

$$f_0 : S(H^Q_0(A), 0) \to A$$

   which induces an isomorphism on H^Q_0.

2. *Suppose A is $(n-1)$-connected for $n \geq 1$. Then there exists a map in $s\mathcal{A}_{\mathbb{F}}$,*

$$f_n : \ S(H^Q_n A, n) \to A,$$

   which is an isomorphism on π_n and H^Q_n.

Proof. (1) Let $\iota : H_0^Q(A) \to I\pi_0 A$ be a choice of splitting for the surjection $I\pi_0 A \to H_0^Q(A)$. By Lemma 3.1, f can be chosen to be the adjoint of the map of simplicial vector spaces $K(H_0^Q(A), 0) \to IA$ induced by ι. By the transitivity sequence, $H_0^Q(A(1)) = 0$ so the fundamental spectral sequence for $A(1)$ converges, by Proposition 2.2 (2), so $A(1)$ is connected. (2) By the Hurewicz theorem, Proposition 2.2 (3), the map $h :\ \pi_n A \to H_n^Q A$ is an isomorphism. Now the adjoint functors

$$S :\ s\mathcal{V} \Longleftrightarrow s\mathcal{A}_{\mathbb{F}} :\ I$$

induce an adjoint pair

$$S :\ Ho(s\mathcal{V}) \Longleftrightarrow Ho(s\mathcal{A}_{\mathbb{F}}) :\ I.$$

Thus we have isomorphisms

$$\begin{aligned} [S(H_n^Q A, n), A] &\cong [K(H_n^Q A, n), IA] \\ &\cong \mathrm{Hom}_{\mathcal{V}}(H_n^Q A, \pi_n IA), \end{aligned}$$

using Lemma 3.1. Choosing f_n to correspond to the inverse of h gives the result. □

Now given A we form the *Postnikov envelopes* as the sequence of cofibrations

$$A(1) \overset{j_2}{\hookrightarrow} A(2) \overset{j_3}{\hookrightarrow} \cdots \overset{j_n}{\hookrightarrow} A(n) \overset{j_{n+1}}{\hookrightarrow} \cdots$$

with the following properties:

(1) $A(1) = \widehat{M(f_0)}$,

(2) for each $n \geq 1$, $A(n)$ is a $(n-1)$-connected and for $s \geq n$,

$$H_s^Q A(n) \cong H_s^Q A.$$

(3) There is a cofibration sequence

$$S(H_n^Q A, n) \to A(n) \overset{j_{n+1}}{\to} A(n+1).$$

The existence of a Postnikov envelopes follows easily from Proposition 2.3, Proposition 3.2, and

Lemma 3.3. *If A is $(n-1)$-connected, for $n \geq 1$, then the cofibre $M(f_n)$ of f_n : $S(H_n^Q A, n) \to A$ is n-connected and satisfies $H_s^Q M(f_n) \cong H_s^Q A$ for $s > n$.*

Proof. This follows from 3.2 and the transitivity sequence

$$H_{s+1}^Q M(f_n) \to H_s^Q S(H_n^Q A, n) \to H_s^Q A \to H_s^Q M(f_n).$$

□

Note: We have been implicitly using the computation

$$H_s^Q S(V, n) = \pi_s QS(V, n) = \pi_s K(V, n) = V$$

for $s = n$ and 0 otherwise. The converse holds as well.

Proposition 3.4. *Let A be connected in $s\mathcal{A}_{\mathbb{F}}$ and suppose $H_s^Q A = 0,\ s \neq n > 0$. Then $A \cong S(H_n^Q A, n)$ in $Ho(s\mathcal{A}_{\mathbb{F}})$.*

Proof. Since A is connected, then A is $(n-1)$-connected by the Hurewicz theorem. By 3.2, $f_n : \; S(H^Q_n A, n) \to A$ is an H^Q_n-isomorphism and hence a weak equivalence by 2.2(2). □

Note. From this proposition, if char$\mathbb{F}$=0 then $S(V,n)$ has simplicial dimension n and $\pi_* S(V,n)$ is free skew-commutative on a basis of V concentrated in degree n. Thus $S(V,n)$ is Q-bounded, for any n, showing that Vanishing Conjecture 1.3 fails in the zero characteristic case.

4. The Homotopy and Homology of n-Extensions

Call an object A in $s\mathcal{A}_\mathbb{F}$ a *simple n-extension* if A is an n-extension in $s\mathcal{A}_\mathbb{F}$ with $\bar{S}_1 = S(V_1, 0)$, V_1 in $\mathcal{V}$. Also, for this section and the next, we define the *simplicial dimension* of A to be

$$s \cdot \dim A = \max\{s |\, H^Q_s(A) \neq 0\}$$

We now proceed to prove:

Theorem 4.1. *Let A be in $s\mathcal{A}_\mathbb{F}$. Then:*

1. *If A is a connected simple n-extension for $n \geq 2$, then, in $Ho(s\mathcal{A}_\mathbb{F})$, we have*
$$A \cong S(H^Q_{n-1}(A), n-1) \otimes S(H^Q_n(A), n).$$
2. *A is a complete intersection if and only if A is a simple 1-extension.*
3. *If A is a complete intersection then $H^Q_s(A) = 0$ for $s \geq 2$ and if $H^Q_*(A)$ is of finite-type then A is bounded.*
4. *If $H^Q_0(A) = 0$ and $H^Q_*(A)$ is of finite-type then $\pi_*(\hat{A})$ is of finite-type.*
5. *The Postnikov envelope $A(1)$ has the following properties:*
    - (a) *If A has finite simplicial dimension, then so does $A(1)$;*
    - (b) *If $H^Q_0(A)$ is finite and $\pi_* A$ is bounded then $\pi_* A(1)$ is Q-bounded.*
    - (c) *If $H^Q_*(A)$ is of finite-type then $H^Q_*(A(1))$ is also of finite-type.*

We begin with

Lemma 4.2. *Let A in $s\mathcal{A}_\mathbb{F}$ be a connected simple n-extension for $n \geq 2$. Then A is an n-extension of the form*

$$S(H^Q_{n-1}A, n-1) \to A \to S(H^Q_n A, n).$$

Proof. Let V_0, V_1 be vector spaces so that there is a cofibration sequence

$$S(V_0, n-1) \to A \to S(V, n).$$

Then the transitivity sequence tells us that $H^Q_s A = 0$, $s \neq n, \; n-1$ and there is an exact sequence

$$0 \to H^Q_n A \to V_1 \to V_0 \to H^Q_{n-1} A \to 0.$$

Thus A is $n-2$ connected and Postnikov tower gives us a cofibration sequence

$$S(H^Q_{n-1}A, n-1) \to A \to A(n-1) = S(H^Q_n A, n).$$ □

Proof of Theorem 4.1 (1). By Lemma 4.2, there is a cofibration sequence

$$S(H^Q_{n-1}, A, n-1) \xrightarrow{i} A \xrightarrow{j} S(H^Q_n A, n),$$

where we can assume A is cofibrant, i is a cofibration, and j is the cofibre. Consider the commuting diagram

$$\begin{tikzcd}
{[S(H^Q_n A, n), A]} \arrow[r, "j_*"] \arrow[d, "\cong"'] & {[S(H^Q_n A, n), S(H^Q_n A, n)]} \arrow[d, "\cong"] \\
{[K(H^Q_n A, n), IA]} \arrow[d, "\cong"'] & {[K(H^Q_n A, n), IS(H^Q_n A, n)]} \arrow[d, "\cong"] \\
\mathrm{Hom}(H^Q_n A, I\pi_n A) \arrow[r, "h_*"] & \mathrm{Hom}(H^Q_n A, H^Q_n A)
\end{tikzcd}$$

Then j will split, up to homotopy, if we can show that $h:\ \pi_n A \to H^Q_n A$ is onto.

By Proposition 2.4(4), there is a commutative diagram

$$\begin{tikzcd}
\pi_n A \arrow[r, "\pi_* j"] \arrow[d, "h"'] & \pi_n S(H^Q_n A, n) \arrow[r, "\tau"] \arrow[d, "\cong"'] & \pi_{n-1} S(H^Q_{n-1}, A, n-1) \arrow[r, "\pi_* i"] \arrow[d, "\cong"] & \pi_{n-1} A \arrow[d, "\cong"] \\
H^Q_n A \arrow[r, "\cong"] & H^Q_n A \arrow[r, "\partial=0"] & H^Q_{n-1} A \arrow[r, "\cong"] & H^Q_{n-1} A
\end{tikzcd}$$

with the rows exact. Thus $\pi_n j$ is onto and, hence, $h:\ \pi_n A \to H^Q_n A$ is onto. □

Lemma 4.3. *Suppose A in $s\mathcal{A}_{\mathbb{F}}$ is regular. Then $S(H^Q_0(A), 0) \cong \hat{A}$ in $Ho(s\mathcal{A}_{\mathbb{F}})$.*

Proof. By the standard transitivity sequence for D_* applied to $\mathbb{F} \to A \to \mathbb{F}$, $D_0(\mathbb{F}|A) = 0$ and $D_{s+1}(\mathbb{F}|A) \cong H^Q_s(A)$, so since A is regular, then $H^Q_s(A) = 0$, $s > 0$. Thus f_0 is an H^Q_*-isomorphism and so $\hat{f}_0$ is a weak equivalence by Proposition 2.3. □

Proof of Theorem 4.1 (2). If A is a complete intersection then it is a 1-extension of the form

$$S_0 \to A \to S_1$$

with S_0 polynomial and $\bar{S}_1$ regular as simplicial augmented algebra. By Lemma 4.3, A is thus a simple 1-extension. The converse is clear. □

Proof of Theorem 4.1 (3). If A is a complete intersection, then $H^Q_s(A) = 0,\ s \geq 2$ follows (2) and the transitivity sequence. Consider now the Eilenberg-Moore spectral sequence

$$E^2_{s,t} = Tor^{S(H^Q_0(A))}_s(\pi_* A, \mathbb{F})_t \Longrightarrow \pi_{s+t} M(f_0)$$

which is a first quadrant homology-type spectral sequence of algebras. Since $H^Q_0(A)$ is finite, $S(H^Q_0(A))$ has finite flat dimension and since, by Proposition 2.3, $Q\pi_* M(f_0) = Q\pi_* A(1) = Q\pi_* S(H^Q_1(A), 1)$ is finite concentrated in degree 1 then $\pi_* M(f_0)$ is bounded and we can conclude, by an induction on $dim_{\mathbb{F}} H^Q_0(A)$, that $\pi_* A$ is bounded. □

Example. Suppose an augmented commutative $\mathbb{F}$-algebra B is a complete intersection. Then there is a complete regular algebra Γ and an ideal I, generated by a regular sequence, so that $\Gamma/I \cong \hat{B}$. As we saw, $\Gamma \cong S(V_0)$, so the condition of regularity on I is equivalent to there being a *projective extension*, that is, (see [7]) an extension

$$\mathbb{F} \to S(V_1) \xrightarrow{i} S(V_0) \to \hat{B} \to \mathbb{F},$$

so that i makes $S(V_0)$ into a projective $S(V_1)$-module. In $Ho(s\mathcal{A}_{\mathbb{F}})$, $\hat{B}$ is equivalent $M(i)$ and so there is a cofibration sequence of the form

$$S(V_0, 0) \to \hat{B} \to S(V_1, 1).$$

Thus, $\hat{B}$, and hence B, is a complete intersection as a simplicial algebra. $\square$

Proof of Theorem 4.1 (4). Since $H_0^Q(A) = 0$, the fundamental spectral sequence

$$E^1_{s,t} = \mathcal{S}_s(H_*^Q(A))_t \Rightarrow \pi_t \hat{A},$$

converges. From the known structure of $\mathcal{S}$ (see e.g. [4]), if V is a finite-dimensional vector space then each $\mathcal{S}_s(V)_t$ is finite and $\mathcal{S}_s(V)_t = 0, s \gg 0$ for each fixed t. The result follows. $\square$

Proof of Theorem 4.1 (5). First, (a) is immediate from the transitivity sequence. For (b), $V = H_0^Q(A)$ is finite and the Eilenberg-Moore spectral sequence has the form

$$E^2_{s,t} = Tor_s^{S(V)}(\pi_* A, \mathbb{F})_t \Longrightarrow \pi_{s+t} M(f_0)$$

Since $S(V)$ has finite flat dimension and $\pi_* A$ is a graded $S(V)$-module then

$$Tor_s^{S(V)}(\pi_* A, \mathbb{F})_t = Tor_s^{S(V)}(\pi_t A, \mathbb{F})$$

vanishes for $s \gg 0$ and vanishes for $t \gg 0$ if $\pi_* A$ is bounded. We conclude $\pi_* M(f_0)$ is bounded and hence $\pi_* A(1)$ is Q-bounded, by Proposition 2.3. Finally, for (c), Lemma 3.3 tells us that $H_s^Q(A) = H_s^Q(A(1))$ for $s \geq 1$. Thus if $H_*^Q(A)$ is of finite-type then $H_*^Q(A(1))$ is of finite-type. $\square$

5. The Poincaré Series of a Simplicial Algebra

For this section, we assume char$\mathbb{F} = p > 0$.

Let A be a connected simplicial augmented commutative $\mathbb{F}$-algebra such that $\pi_* A$ is of finite-type. We define its *Poincaré series* by

$$\vartheta(A, t) = \sum_{n \geq 0} (dim_{\mathbb{F}} \pi_n A) t^n.$$

If V is a finite-dimensional vector space and $n > 0$ we write

$$\vartheta(V, n, t) = \vartheta(S(V, n), t).$$

Given power series $f(t) = \sum a_i t^i$ and $g(t) = \sum b_i t^i$ we define the relation $f(t) \leq g(t)$ provided $a_i \leq b_i$ for each $i \geq 0$.

Lemma 5.1. *Given a cofibration sequence*

$$A \to B \to C$$

of connected objects in $\mathcal{A}_{\mathbb{F}}$ *with finite-type homotopy groups, then*

$$\vartheta(B,t) \leq \vartheta(A,t)\vartheta(C,t)$$

which is an equality if the sequence is split.

Proof. From the Serre spectral sequence

$$E^2_{s,t} = \pi_s(C \otimes \pi_t A) \Longrightarrow \pi_{s+t}B$$

we have

$$\vartheta(A,t)\vartheta(C,t) = \sum_n (\sum_{s+t=n} dim_{\mathbb{F}} E^2_{s,t})t^n \geq \vartheta(B,t).$$

If the cofibration sequence is split then the spectral sequence collapses, giving an equality. □

If Π is a finitely-generated abelian group and $n > 0$ let

$$\vartheta(\Pi,n,t) = \sum_s (dim_{\mathbb{F}} H_s(K(\Pi,n);\mathbb{F}))t^s.$$

Lemma 5.2. *Let* V *be a finite-dimensional vector space and* Π *a free abelian group of the same dimension. Then for any* $n > 0$

$$\vartheta(V,n,t) = \vartheta(\Pi,n,t).$$

Proof. As shown in [5], there is a weak equivalence of simplicial vector spaces

$$S(V,n) \to \mathbb{F}[K(\Pi,n)]$$

which gives us the desired result. □

Proposition 5.3. *Given a finite-dimensional vector space* V *and any* $n > 0$ *the Poincaré series* $\vartheta(V,n,t)$ *converges in the open unit disc.*

Proof. This follows from Lemma 5.2 and the results of J.P. Serre in [14] and Y. Umeda in [16]. □

Now given two power series $f(t)$ and $g(t)$ we say $f(t) \sim g(t)$ provided $lim_{t\to\infty} f(t)/g(t) = 1$. Given a Poincaré series $\vartheta(V,n,t)$, for a finite-dimensional $\mathbb{F}$-vector space V and $n > 0$, let

$$\varphi(V,n,t) = log_p \vartheta(V,n,1-p^{-t}).$$

Proposition 5.4. *For* V *an* $\mathbb{F}$*-vector space of finite dimension* q *and* $n > 0$ *then* $\varphi(V,n,t)$ *converges on the real line and*

$$\varphi(V,n,t) \sim qt^{n-1}/(n-1)!.$$

Proof. This follows from Lemma 5.2 and Théorème 9b in [14], for char$\mathbb{F} = 2$, and its generalization in [16]. □

A major step in proving Theorem 1.4 will be accomplished with

Theorem 5.5. *Let A be a connected finite simplicial augmented commutative $\mathbb{F}$-algebra. Then if A is Q-bounded we have $A \cong S(H_1^Q(A), 1)$ in $Ho(s\mathcal{A}_{\mathbb{F}})$.*

Proof. By Theorem 4.1 (4) and Proposition 2.3, $\pi_* A$ is of finite-type and hence, as it is also Q-bounded, bounded as well. Let $n = s \cdot \dim A$. We must show that $n = 1$.

Consider the Postnikov envelope

$$S(H_{s-1}^Q(A), s-1) \to A(s-1) \to A(s)$$

for each s. From the theory of cofibration sequences (see section I.3 of [11]) the above sequence extends to a cofibration sequence

$$A(s-1) \to A(s) \to S(H_{s-1}^Q(A), s).$$

Thus, by Lemma 5.1, we have

$$\vartheta(A(s), t) \leq \vartheta(A(s-1), t)\vartheta(H_{s-1}^Q(A), s, t).$$

Starting at $s = n-1$ and iterating this relation, we arrive at the inequality

$$\vartheta(A(n-1), t) \leq \vartheta(A, t) \prod_{s=1}^{n-2} \vartheta(H_s^Q(A), s+1).$$

Now, $A(n) \cong S(H_n^Q(A), n)$ by Proposition 3.4, but, by Theorem 4.1 (1) and Lemma 5.1, we have

$$\vartheta(A(n-1), t) = \vartheta(H_{n-1}^Q(A), n-1, t)\vartheta(H_n^Q(A), n, t).$$

Since $\pi_*(A)$ is of finite-type and bounded then there exists a $D > p$ such that $\vartheta(A, t) \leq D$, in the open unit disc. Combining, we have

$$\vartheta(H_{n-1}^Q(A), n-1, t)\vartheta(H_n^Q(A), n, t) \leq D \prod_{s=1}^{n-2} \vartheta(H_s^Q(A), s+1).$$

Applying a change of variables and log_p to the above inequality, we get

$$\varphi(H_{n-1}^Q(A), n-1, t) + \varphi(H_n^Q(A), n, t) \leq d + \sum_{s=1}^{n-2} \varphi(H_s^Q(A), s+1).$$

By Proposition 5.4, there is a polynomial $f(t)$ of degree $n-2$, non-negative integer a, and positive integers b and d such that

$$at^{n-2} + bt^{n-1} \leq d + f(t)$$

which is clearly false for $n > 1$. Thus $n = 1$. The rest of the proof follows from Proposition 3.4. □

Proof of Theorem 1.4. The "if" part is Theorem 4.1 (3). We thus concentrate on the "only if" part. We are given a finite simplicial augmented commutative $\mathbb{F}$-algebra A

such that π_*A is bounded. By Theorem 4.1 (4) and (5), $A(1)$ is connected, Q-bounded, finite, and $\pi_*A(1)$ is of finite-type. Thus $A(1)$ is bounded as well and we conclude $A(1) \cong S(H_1^Q(A), 1)$ by Theorem 5.5. $\square$

References

[1] M. André, *Homologie des Algèbres Commutatives*, Die Grundlehren der Mathematischen Wissenschaften 206, Springer-Verlag, 1974.

[2] L. Avramov, Local rings of finite simplicial dimension, Bull. A.M.S. (New Ser.) 10(1984), 289–291.

[3] ———, Locally complete intersection homomorphisms and a conjecture of Quillen on the vanishing of cotangent homology, preprint, Purdue University 1997.

[4] A.K. Bousfield, Operations on derived functors of non-additive functors, manuscript, Brandeis University 1967.

[5] H. Cartan, *Algèbres d'Eilenberg-MacLane et homotopie*, Exposés 2 à 11, Sém. H. Cartan, Éc. Normale Sup. (1954–1955), Sectétariat Math., Paris, 1956; [reprinted in:] *Œvres*, vol. III, Springer, Berlin, 1979; pp. 1309–1394.

[6] A. Dold, Homology of symmetric products and other functors of complexes, Ann. of Math. (2) 68 (1958), 40-80.

[7] P. Goerss, On the André-Quillen cohomology of commutative $\mathbb{F}_2$-algebras, Astérique 186(1990).

[8] ———, A Hilton-Milnor theorem for categories of simplicial algebras, Amer. J. Math,. 111(1989), 927–971.

[9] S. MacLane, *Homology*, Classics in Mathematics, Springer-Verlag, 1995.

[10] H. Miller, The Sullivan conjecture on maps from classifying spaces, Annals of Math. 120(1984), 39–87.

[11] D. Quillen, *Homotopical Algebra*, Lecture Notes in Mathematics 43, Springer-Verlag, 1967.

[12] ———, On the (co)homology of commutative rings, Proc. Symp. Pure Math. 17(1970), 65–87.

[13] ———, On the homology of commutative rings, Mimeographed Notes, M.I.T.

[14] J.-P. Serre, Cohomologie modulo 2 des espaces d'Eilenberg-MacLane, Comment. Math. Helv. 27 (1953), 198-231

[15] J.M. Turner, Completions and localizations of simplicial commutative algebras, forthcoming.

[16] Y. Umeda, A remark on a theorem of J.-P. Serre, Proc. Japan. Acad. 35 (1958), 563-568.

Department of Mathematics, College of the Holy Cross, One College Street, Worcester, MA 01610-2395

E-mail address: `jmturner@math.holycross.edu`